

PATIENT INFORMATION CRESTOR® (Kres-tor) (rosuvastatin) tablets, for oral use
Read this Patient Information carefully before you start taking CRESTOR and each time you get a refill. If you have any questions about CRESTOR, ask your healthcare provider. Only your healthcare provider can determine if CRESTOR is right for you.
What is CRESTOR? CRESTOR is a prescription medicine that contains a cholesterol-lowering medicine called rosuvastatin. - CRESTOR is used: - to reduce the risk of major adverse cardiovascular (CV) events, such as death from cardiovascular disease, heart attack, stroke, or the need for procedures to improve blood flow to the heart called arterial revascularization, in adults who do not have known heart disease but do have certain additional risk factors. - along with diet to: - lower the level of low-density lipoprotein (LDL-C) cholesterol or “bad” cholesterol in adults with primary hyperlipidemia. - slow the buildup of fatty deposits (plaque) in the walls of blood vessels. - treat adults and children 8 years of age and older with high blood cholesterol due to heterozygous familial hypercholesterolemia (HeFH) (an inherited condition that causes high levels of LDL-C). - along with other cholesterol lowering treatments or alone if such treatments are unavailable in adults and children 7 years of age and older with homozygous familial hypercholesterolemia (HoFH) (an inherited condition that causes high levels of LDL-C). - along with diet for the treatment of adults with: - primary dysbetalipoproteinemia (an inherited condition that causes high levels of cholesterol and fat). - hypertriglyceridemia. It is not known if CRESTOR is safe and effective in children younger than 8 years of age with HeFH or children younger than 7 years of age with HoFH or in children with other types of hyperlipidemias (other than HeFH or HoFH).
Do not take CRESTOR if you: - have liver problems. - are allergic to rosuvastatin or any of the ingredients in CRESTOR. See the end of this leaflet for a complete list of ingredients in CRESTOR.
Before you take CRESTOR, tell your healthcare provider about all of your medical conditions, including if you: - have unexplained muscle aches or weakness. - have or have had kidney problems. - have or have had liver problems. - drink more than 2 glasses of alcohol daily. - have thyroid problems. - are of Asian descent. - are pregnant or think you may be pregnant, or are planning to become pregnant. If you become pregnant while taking CRESTOR, call your healthcare provider right away to discuss your CRESTOR treatment.

- are breastfeeding. CRESTOR can pass into your breast milk and may harm your baby. Talk to your healthcare provider about the best way to feed your baby if you take CRESTOR. Do not breastfeed while taking CRESTOR.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Tell your healthcare provider who prescribes CRESTOR if another healthcare provider increases the dose of another medicine you are taking. CRESTOR may affect the way other medicines work, and other medicines may affect how CRESTOR works.

Especially tell your healthcare provider if you take:

- coumarin anticoagulants (medicines that prevent blood clots, such as warfarin)
- antacids (medicines you take for heartburn that contain aluminum and magnesium hydroxide)

Taking CRESTOR with certain medicines may increase the risk of muscle problems.

Especially tell your healthcare provider if you take:

- cyclosporine (a medicine for your immune system)
- teriflunomide (a medicine used to treat relapsing remitting multiple sclerosis)
- enasidenib (a medicine used to treat acute myeloid leukemia)
- capmatinib (a medicine for the treatment of non-small cell lung cancer)
- fostamatinib (a medicine used to treat low platelet counts)
- febuxostat (a medicine used to treat and prevent high blood levels of uric acid)
- gemfibrozil (a fibric acid medicine for lowering cholesterol)
- tafamidis [used to treat cardiomyopathy (enlarged and thickened heart muscle)]
- anti-viral medicines including certain HIV or hepatitis C virus drugs such as:
 - lopinavir, ritonavir, fosamprenavir, tipranavir, atazanavir, simeprevir
 - combination of
 - sofosbuvir/velpatasvir/voxilaprevir
 - dasabuvir/ombitasvir/paritaprevir/ritonavir
 - elbasvir/grazoprevir
 - sofosbuvir/velpatasvir
 - glecaprevir/pibrentasvir **and**
 - all other combinations with ledipasvir including ledipasvir/sofosbuvir
- darolutamide (a medicine for the treatment of prostate cancer)
- regorafenib (a medicine used to treat cancer of the colon and rectum)
- fibric acid derivatives (such as fenofibrate)
- ticagrelor (helps reduce the chance of a blood clot formation that can block a blood vessel)
- niacin or nicotinic acid
- colchicine (a medicine used to treat gout)

Ask your healthcare provider or pharmacist for a list of these medicines if you are not sure. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get new medicine.

How should I take CRESTOR?

- Take CRESTOR exactly as your healthcare provider tells you to take it.
- Take CRESTOR, by mouth, 1 time each day, with or without food. Swallow the tablet whole.
- CRESTOR can be taken at any time of day, with or without food.
- **Do not** change your dose or stop CRESTOR without talking to your healthcare provider, even if you are feeling well.

- Your healthcare provider may do blood tests to check your cholesterol levels before and during your treatment with CRESTOR. Your healthcare provider may change your dose of CRESTOR if needed.
- While taking CRESTOR, continue to follow your cholesterol-lowering diet and to exercise as your healthcare provider told you to.
- If you take a medicine called an antacid that contains a combination of aluminum and magnesium hydroxide, take CRESTOR at least 2 hours before you take the antacid.
- If you miss a dose of CRESTOR, take your next dose at your normal scheduled time. **Do not take** an extra dose of CRESTOR.
- In case of an overdose, get medical help or contact a live Poison Center expert right away at 1-800-222-1222. Advice is also available online at poisonhelp.org.

What are the possible side effects of CRESTOR?

CRESTOR may cause serious side effects, including:

- **Muscle pain, tenderness and weakness (myopathy).** Muscle problems, including muscle breakdown, can be serious in some people and rarely cause kidney damage that can lead to death. Tell your healthcare provider right away if:
 - you have unexplained muscle pain, tenderness, or weakness, especially if you have a fever or feel more tired than usual, while you take CRESTOR.
 - you have muscle problems that do not go away even after your healthcare provider has told you to stop taking CRESTOR. Your healthcare provider may do further tests to diagnose the cause of your muscle problems.

Your chances of getting muscle problems are higher if you:

- are taking certain other medicines while you take CRESTOR (see “Especially tell your healthcare provider if you take”)
- are 65 years of age or older
- are of Asian descent
- have thyroid problems (hypothyroidism) that are not controlled
- have kidney problems
- are taking higher doses of CRESTOR
- **Liver problems.** Your healthcare provider may do blood tests to check your liver before you start taking CRESTOR and if you have symptoms of liver problems while you take CRESTOR. Call your healthcare provider right away if you have any of the following symptoms of liver problems:
 - feel unusually tired or weak
 - loss of appetite
 - upper belly pain
 - dark urine
 - yellowing of your skin or the whites of your eyes
- **Protein and blood in the urine.** CRESTOR may cause you to have protein and blood in your urine. If you develop protein or blood in your urine, your healthcare provider may decrease your dose of CRESTOR.
- **Increase in blood sugar (glucose) levels.** CRESTOR may cause an increase in your blood sugar levels.

The most common side effects may include headache, nausea, muscle aches and pains, weakness, and constipation.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store CRESTOR?

- Store CRESTOR at room temperature, between 68°F to 77°F (20°C to 25°C) and in a dry place.

Keep CRESTOR and all medicines out of the reach of children.

General Information about the safe and effective use of CRESTOR

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use CRESTOR for a condition for which it was not prescribed. Do not give CRESTOR to other people, even if they have the same medical condition you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about CRESTOR that is written for health professionals.

What are the Ingredients in CRESTOR?

Active Ingredient: rosuvastatin as rosuvastatin calcium

Inactive Ingredients: crospovidone NF, hypromellose NF, lactose monohydrate NF, magnesium stearate NF, microcrystalline cellulose NF, red ferric oxide NF, titanium dioxide USP, triacetin NF, tribasic calcium phosphate NF and yellow ferric oxide.

CRESTOR® is a trademark of the AstraZeneca group of companies.

© AstraZeneca 2024

Licensed from SHIONOGI & CO., LTD., Osaka, Japan

Distributed by:

AstraZeneca Pharmaceuticals LP

Wilmington, DE 19850

For more information, go to the CRESTOR website at www.crestor.com or call 1-800-CRESTOR

This Patient Information has been approved by the U.S. Food and Drug Administration

Revised 07/2024